Taiwan: Obesity Health System





Health system summary

Taiwan has a universal national health insurance programme that is mandatory for all its citizens (and for internationals residing in Taiwan for longer than six months). This single-payer, compulsory social insurance covers most of the population - 99.9% of the population in 2016 - and is mostly financed through payroll-based premiums.² The rest of the funding comes from government funding and out of pocket payments. Private health insurance does not tend to cover services provided by the public health insurance and therefore does not ensure faster access to services and specialists. Overall, out of pocket expenditure is estimated to be approximately 26% of total financing.³



WØRLD ØBESITY

www.worldobesity.org

Taiwan: Obesity Health System

Obesity prevalence

6.4%	6.1%
Women	Men
3.6%	8.2%
Girls	Boys

Overweight prevalence

UNK% Women	UNK% Men
13.0%	18.6%
Girls	Boys

Key prevention policies

- ⇒ Mandatory nutrient lists on packaged foods
- ⇒ Restrictions on food advertising and promotion to children on broadcast

Summary of stakeholder feedback*

Stakeholders praised Taiwan's universal health insurance system that covers nearly all of its population. It was said that people can generally enter the health system freely, but for obesity there was disagreement between stakeholders about when one with obesity would be picked up by the system. Responses ranged from when people had a BMI >27 kg/m² to >40 kg/m² and one pointed out that patients were more likely to seek treatment is they had higher education and/or a higher income.

Stakeholders reported that Taiwan's national health insurance generally does not pay for the medical management of obesity. Consequently, obesity treatment is largely covered by out-of-pocket expenditure. This reflects the poor investment into obesity, with obesity management only just starting to be recognised in the health system. It was said that the national health insurance only covered the treatment of severe obesity with bariatric surgery when BMI was >40 kg/m² or between 35-39.9 kg/m² with comorbidities. As a result, most patients living with obesity are not adequately treated. It was reported that people tended to leave the system because of this limited coverage or because they were not referred onto specialist treatment.

The stakeholders noted that there are clinical guidelines for the prevention and management of obesity in both adults and children. However, these are not well implemented yet and there is a need to educate more health professionals and providers about obesity. Training for health professionals was considered limited, with there being a lack of suitably qualified professionals in both urban and rural areas. Stakeholders specifically reported a lack of psychiatrists and psychologists.

*Based on interviews/survey returns from 3 stakeholders



Perceived barriers to treatment



References

3.

- 1. <u>https://blogs.worldbank.org/opendata/new-country-</u> <u>classifications-income-level-2019-2020</u> (last accessed 29.08.19)
- 2. <u>https://international.commonwealthfund.org/countries/</u> taiwan/ (last accessed 14.05.20)
 - http://gnhe.org/blog/wp-content/uploads/2015/05/GNHE-UHC-assessment_Taiwan-1.pdf (last accessed 13.11.2018)

Last updated June 2020